

AN ASPIRIN DIRECTORY

These common, brand-name prescription and over the counter drugs contain aspirin. It's necessary to stop using them at **least two weeks prior to surgery** and for one week after. They can cause bleeding and bruising. This list does not include any generic preparations sold under store brands or other names.

Alka-Seltzer tablets	Empirin
Alka-Seltzer Plus cold medicine	Empirin with Codeine
Anacin capsules and tablets	Emprazil tablets
APC tablets	Emprazil-C tablets En Tab
APC with Bultalbital tablets	Excedrin
APC with Codeine Tabloid brand	Extra Strength Bufferin
Anacin Maximum Strength	Fiorinal
Arthritis Pain Formula (Anacin)	Fiorinal with Codeine
Arthritis Strength Bufferin	4-Way Cold Tablets
Ascriptin	Goody's Headache Powders
Aspergum	Midol
Aspirin suppositories	Momentum Muscular Backache Formula
Bayer aspirin	Norgesic
Bayer Children's Chewable Aspirin	Norgesic Forte
Bayer Children's Cold Tablets	Norwich Aspirin
Bayer Timed-Released Aspirin	Pabirin buffered tablets
Buff-A Comp tablets	Panalgesic
Bufferin	Percodan and Percodan-Demi tablets
Cama Inlay-Tabs	Quiet World Analgesic/Sleeping Aid
Cetased, Improved	Robaxisal tablets
Gongesprin	SK-65 Compound
Coricidin D decongestant tablets	St. Joseph Aspirin for Children
Coricidin for children	Sine-Off Sinus Medicine
Coricidin Medilets tablets for children	Supac
Coricidin tablets	Synalgos capsules
Darvon with A.S.A.	Synalgos-DC capsules
Darvon-N with A.S.A.	Triaminicin tablets
Gemnisyn	Vanquish
Dristan decongestant	Verin
Ecotrin tablets	Viro-Med tablets
	Zorprin